



Gastroenterology - Hepatology - Diagnostic & Therapeutic Endoscopy

A/Prof David van der Poorten - Dr Roslyn Vongsuvanh - Dr Way Siow Prof Golo Ahlenstiel - Dr Rishi Sud - Dr Wai See Ma

THREE DAYS BEFORE PROCEDURE

Lakeview Private Hospital Suite 4, 17-19 Solent Circuit Norwest NSW 2153 Phone: 8711 0160 - Fax: 8624 4769

INSTRUCTIONS FOR AFTERNOON COLONOSCOPY - BOWEL PREPARATION WITH PREPKIT ORANGE

THIS KIT CONTAINS: 1 Glycoprep Orange 70g sachet 2 Picoprep Orange 20g sachets

NOTE: The success of your examination depends on the bowel being as clear as possible, otherwise the examination may need to be postponed and the preparation repeated. This preparation will cause multiple bowel movements and often works quickly. Remain within easy reach of a toilet.

Stop all iron containing medicine 7 days prior to the procedure. Continue with your regular medications. Special instructions are needed if you are diabetic, or on Clopidogrel, warfarin or other blood thinners.

ADVERSE REACTIONS: The most frequent reactions are nausea and bloating. These can be reduced by slowing the rate of intake. Less frequent adverse reactions include vomiting and rectal irritation (Vaseline may assist). These adverse reactions are temporary and generally disappear rapidly.

7pm T	ake 2 Senokot tablets (brown)
Commence	B BEFORE PROCEDURE e modified residue fibre diet – refer to attached sheet. ake 2 Senokot tablets (brown)
Drink only	CLEAR FLUIDS for breakfast, lunch and dinner. FOODS OR MILK PRODUCTS ALLOWED.
Gatorade,	OCLEAR FLUIDS: (drink a mix) Powerade, other electrolyte sports drinks. Strained fruit juices, water, clear broth, Bonox, tea, coffee ellies (not red or purple), cordials (not red or purple), clear ice blocks.
6РМ	Dissolve the entire contents of 1 sachet of Picoprep Orange in 250ml of warm water. If preferred chill in fridge. Drink contents of glass followed by 1 glass of water.
Continue drinking clear fluids as above at least 2-3 glasses /hour/the more the better.	
9PM	Dissolve the entire contents of 1 sachet of Glycoprep Orange in 1L of water. If preferred chill in fridge prior. Drink mixture slowly over 1 hour (one glass every 15mins).
Continue drinking clear fluids as above at least 2-3 glasses /hour/the more the better.	
DAY OF PROCEDURE	
8AM	Dissolve the entire contents of 1 sachet of Picoprep Orange in 250ml of warm water. If preferred

chill in fridge. Drink contents of glass followed by 1 glass of water.

Continue drinking clear fluids up until 2 hours prior to your procedure time, then FAST (no food/drink).

Please note that you are NOT allowed to drive home after your procedure.

MODIFIED RESIDUE DIET

The diet aims to avoid foods which are incompletely digested or which may act as irritants to the bowel. Example: Fried foods, highly seasoned food and foods which contain a high fibre diet content.

CEREALS:

Avoid wholegrain cereals (WEET-BIX), All Bran, brown/wholemeal breads, brown rice, wholemeal pastas, wholemeal biscuits or those containing nuts or fruit.

Allowed:

Cornflakes, Rice Bubbles, Strained rolled oats, Semolina, white bread, white rice, white pasta, plain biscuits (Arrowroot, Sao, Milk Coffee).

VEGETABLES:

Avoid all varieties.

FRUIT:

Avoid all fresh, canned, stewed or dried fruit.

Allowed:

Strained fruit juices.

MEAT, FISH, POULTRY, EGGS:

Avoid red meat (lamb, beef, veal, pork, bacon, ham), made up meats (devon, rissoles, curries, highly seasoned sausages or fatty casseroles), fried fish, canned fish, shellfish, fried eggs.

Allowed:

Chicken with no skin, rabbit, fish (boiled, steamed or grilled), eggs (boiled, poached, scrambled, plain omlets).

MILK AND CHEESE:

Avoid strong cheeses.

Allowed:

300ml milk (whole or skim), 50g mild cheese.

FATS:

Avoid cooking and salad oils.

Allowed.

30g or 1 ½ tablespoons butters or margarine or 60ml of cream (3 tablespoons).

BEVERAGES:

Avoid alcohol.

Allowed:

Tea, coffee, milk from allowance, soft drinks, cordials, strained cream soups, broth, Bonox.

DESSERTS:

Avoid pastries, puddings containing fruit, dried nuts or fruits.

Allowed:

Milk puddings, jellies, plain sponge.

MISCELLANEOUS:

Avoid jam, marmalade, pickles, chutney, sauces, mayonnaise, nuts, coconut, fudge, chocolate, herbs and spices. *Allowed:*

Sugar, honey, boiled lollies, barley sugar, Marmite, Vegemite, Bonox, salt.